

In Vitro Comparative Study of Osteogenic Differentiation Ability between Adipose and Bone Marrow Mesenchymal Stem Cell Applied to Bovine Demineralized Bone Matrix

by Dwikora Novembri

Submission date: 21-Nov-2018 02:45PM (UTC+0800)

Submission ID: 1043084882

File name: Dr_FER-DNU-LUK-MED_JBBBE_2018_2_2.pdf (3.38M)

Word count: 3684

Character count: 19975

In Vitro Comparative Study of Osteogenic Differentiation Ability between Adipose and Bone Marrow Mesenchymal Stem Cell Applied to Bovine Demineralized Bone Matrix

Ferdiansyah Mahyudin^{1,a,*}, Dwikora Novembri Utomo^{1,b}, Mouli Edward^{1,c},
Lukas Widhiyanto^{1,d}, Christopher Anthony Simanjuntak^{1,e},
Hudaya Nikmatullah^{1,f}

¹Department of Orthopedic and Traumatology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Hospital, Indonesia

^aferdyortho@gmail.com, ^bdwikora-novembri-u@fk.unair.ac.id, ^cmouli-edward@fk.unair.ac.id,
^dlukas-w@fk.unair.ac.id, ^ecas.orthosby@gmail.com, ^fhudaya.nikmatullah@gmail.com

*Corresponding author: ferdyortho@gmail.com

Keywords: bone defect, demineralized bone matrix, mesenchymal stem cell

Abstract. Ideal bone graft must possess the desirable trait such as osteoconductive, osteoinductive and osteogenesis. Demineralized Bone Matrix (DBM) provides both osteoconductive and osteoinductive trait. Referring to the tissue engineering principle, the addition of mesenchymal stem cell would add the osteogenic trait to this procedure. The design of this study is experimental using Bovine DBM. Bone Marrow Mesenchymal Stem Cell (BMSCs) and Adipose Mesenchymal Stem Cells (ASCs) were taken from New Zealand white rabbit. There are two groups of treatment, divided into DBM implanted with BMSCs and DBM implanted with ASCs. Each BMSCs and ASCs groups is incubated in the normal and osteogenic culture plate. Evaluation is performed by counting the osteoblast and immunohistochemistry stain using Alkaline Phosphate and Osteocalcin. After 4 weeks of incubation, we found that the osteoblast count in BMSCs groups is higher compared to the ASCs groups in both culture condition ($p < 0.01$) along with Alkaline Phosphate staining ($p < 0.05$), while the Osteocalcin staining showed insignificant differences ($p > 0.05$). This study revealed that xenogenic bovine DBM can act as the potential osteoinductive scaffold for the MSCs to differentiate. The tissue engineering application by combining MSCs and Bovine DBM can be considered as an alternative in managing bone defect cases.

1. Introduction

Bone defects are one of the major problems in orthopedic cases. It can occur in both trauma and non trauma cases. In bone defects, discontinuities of bone integrity can be caused by high-energy trauma, infection, tumor resection surgery, revision surgery, developmental deformities, and congenital malformation disorders. In this case, the size of the bone defect is too large and is included in the critically sized defect so it will not undergo spontaneous regeneration and require an interventional operation [1,2]. There are 500,000 surgeries that require bone grafting in the United States within a year with a total cost of 300 million USD in 1999 [3]. For comparison, the use of bone graft for orthopedic surgery at Dr. Soetomo General Hospital Surabaya has quadrupled from 1997 to 2001 [4]. Bone defect treatment requires the bone graft to fill the gap and autologous bone graft is still a gold standard in the treatment of bone defect cases [5]. However, postoperative pain in bone donor source areas is of concern in the use of autologous bone graft besides its limitations of the bone sources [6]. Other alternatives for the graft substitutes are allogenic bone graft and animals xenogenic bone graft. Allograft is physiologically more resembling the original bone and may have osteoinductive properties but the donor availability is also limited and disease transmission still be concerned. Whereas xenograft, owing to the abundance of donors, may be less expensive and more readily available even though its properties are limited to osteoconductive alone [7].

With the rapid progress in the concept of tissue engineering, the use of stem cells that originally are exogenous cells in bone tissue engineering serves osteogenic capacity in stimulating bone healing [8,9]. Together with the development of scaffolding techniques by performing demineralization so cellular components that potentially induce immune response can be eliminated, thus providing potential scaffold as a microenvironment of the stem cell [8].

Considering this, bovine Demineralized Bone Matrix (DBM) which is a xenogenic bone graft product can be rated as the alternative bone graft substitutes, because its osteoconductive and osteoinductive properties can serve as a scaffold to support stem cell proliferation and differentiation thus provide osteogenic capacity [9]. Besides its economically cost, bovine DBM has been shown to have higher bone healing promotion capacities than the other type of bone graft and has lower immunogenicity and rejection reactions to surrounding tissue [10]. This current study will test and compare the ability of bovine DBM as scaffold and growth factor provider to stimulate differentiation of bone marrow and adipose mesenchymal stem cells to enhance its function as a bone graft.

2. Materials and Methods

This study was an in vitro laboratory study conducted at the Institute of Tropical Disease (ITD) Airlangga University and Tissue Bank of Dr. Soetomo General Hospital in Surabaya, Indonesia. The experimental unit was BMSCs and ASCs isolated from a healthy male New Zealand white rabbit 8 months old and weighing 3.0 kg approved by Airlangga University and Dr. Soetomo General Hospital research committee. In this study, bovine DBM added to isolated and replicated BMSCs and ASCs and divided into 4 groups (Figure 1).




Figure 1. Research design

2.1 Preparation of bovine demineralized bone matrix

Demineralized bone matrix, prepared from the femoral condyle of the 2-year-old local cow, were collected from the local slaughterhouse. All bones were collected aseptically, and the soft tissues were removed before storage at -70°C. The bones were later cleared of fascia and cut into 1-cm pieces with a bone saw under saline (0.9% NaCl) solution lavage. Bone pieces were stored at -70°C until further use. The pieces were then thawed in 95% ethanol for 15 minutes and air dried. All bones were milled to 10 mm pieces. The pieces were then decalcified in 0.6 mol/l HCl at 4°C for 8 days under constant agitation. After demineralization, all bone pieces were rinsed in sterile water and placed in phosphate buffer overnight. The bone pieces were then rinsed and the pH was adjusted to 7.3. They were placed in ethanol, the ethanol was allowed to evaporate overnight, and the pieces were packaged aseptically and stored at 4°C.

2.2 Isolation of bone marrow mesenchymal stem cells

Animals were anesthetized with ketamine (40 mg/kg, IM) and xylazine (5 mg/kg, IM). Isolation of MSCs from bone marrow performed aseptically by cleaning the skin of rabbit femur. The muscle and connective tissue were removed from the femur by scraping the diaphysis of the bone clean

then pulling the tissue toward the ends of the bone. A 27-gauge needle was inserted and flushed with Dulbecco's Modified Eagle's Medium (DMEM) and collect in a 15mL tube. The cell suspension was filtered through a 70- μ m filter mesh. Bone marrow cells were cultured in DMEM+10% FBS+1% antibiotic-antimycotic solution tissue culture flask and incubated at 37°C with 5% CO₂. For osteogenic culture, 10 mmol dexamethasone, 10 mmol β -glycerol-phosphate, 0.05 mmol 2-phosphate ascorbic acid, 100 U/mL penicillin/streptomycin, and 10 mmol 1,25-dihydroxy vitamin D3 were added to culture plate.

2.3 Isolation of adipose mesenchymal stem cells

Isolation of MSCs from adipose tissue was performed by subcutaneous fat pad liposuction under sterile conditions and washed with phosphate-buffered saline (PBS) containing 1% antibiotic-antimycotic solution. The washing step was repeated until all blood vessels and connective tissues appeared to have been liberated. Adipose tissue then minced into small pieces and digested in 0.1% type 1 collagenase at 37°C with shaking for 2 hours and added 5mL DMEM containing 10% fetal bovine serum. For further disintegration of tissue aggregates, the sample was pipetted up and down several times. The cell suspension was filtered through 100 μ m filter for the removal of the solid aggregates. The sample was subsequently centrifuged at 2000 rpm for 5 minutes at room temperature to complete the separation of the stromal cells from the adipocytes and the supernatant removed without disturbing the cells. The pellet was resuspended in 1 ml of lysis buffer to lyse red blood cells, incubated for 10 minutes, washed with 10 ml of PBS+1% antibiotic-antimycotic mixture and centrifuged at 2000 rpm for 5 minutes. The supernatant was removed and the cell pellet was resuspended in complete medium (DMEM with 20% fetal bovine serum (FBS) and 1% antibiotic-antimycotic solution) in a 25 cm² culture flask and maintained in an incubator supplied with the humidified atmosphere of 5% CO₂ at 37°C.

2.4 Cell cultivation

After one day, non-adherent cells were removed by two to three washes with PBS and adherent cells further cultured in complete medium. The medium was changed every 3 days until the monolayer of adherent cells reached 70-80% confluence. Cell passaging was performed using trypsin-EDTA solution 0.25%. Cell cultivation was performed up to the 3rd passage.

2.5 Osteogenic differentiation

Passage 3 MSCs were harvested by trypsin digestion as described above. The cells were counted and seeded at a density of 5×10^4 per well in a 9-well plate each group. The medium was changed twice per week for 2-3 weeks. The differentiation potential for osteogenesis was observed by the number of osteoblasts, alkaline phosphatase (ALP), and osteocalcin (OC). The number of osteoblasts assessed by hematoxylin-eosin staining, osteogenic marker gene of alkaline phosphatase and osteocalcin assessed by immunohistochemical staining under immunofluorescence microscopy. The Immunoreactive Score (IRS) by Remmele is a scale index is used to quantify the immunohistochemistry findings resulted by multiplication of immunoreactive cell percentage scores with color intensity scores on immunoreactive cells (Table 1). The data of each microscopic examinations were observed in 5 fields of view.

Table 1. The semi-quantitative scale of the IRS (IRS result = AxB)

A	B
Score 0: no positive cells	Score 0: no color reactions
Score 1: positive cells less than 10%	Score 1: low color intensity
Score 2: positive cells ranging from 11% - 50%	Score 2: medium color intensity
Score 3: positive cells between from 51% - 80%	Score 3: strong color intensity
Score 4: positive cells between from more than 80%	

3. Results and Discussion

The MSCs cultures were observed by using an inverted light microscope. Adherence of spindle-shaped cells to culture plastic flask was observed after 1 day of culture for both BMSCs and ASCs. Primary cultures reached 70-80% confluence faster in BMSCs than for ASCs (Figure 2).




Figure 2. (A) Bone marrow stem cells culture, (B) Adipose stem cells culture

Table 2 and Figure 3 reveal the mean total number of osteoblast cells in the BMSCs group added with bovine DBM and cultured at normal condition was 7.67 ± 2.00 cells per 5 fields of view, higher than the osteoblast cell number in the ASCs group (1.44 ± 1.24) and the difference is statistically significant ($p = 0.000$). Similar significant result found for the mean of osteoblast number cells in the osteogenic culture, BMSCs group was 120.67 ± 28.54 compared to ASCs group 73.00 ± 42.32 cells number per 5 fields of view ($p = 0.013$).

Table 2. Osteogenic differentiation of mesenchymal stem cells

	BMSCs (Mean ± SD)		ASCs (Mean ± SD)	
	Culture Condition			
	Normal	Osteogenic	Normal	Osteogenic
Osteoblast Count (Cell number/ 5 FV)	7.67±2.00	120.67±28.54	1.44±1.24	73.00±42.32
ALP Expression (IRS Score)	3.20±0.63	3.84±0.53	1.12±0.36	1.16±0.33
Osteocalcin Expression (IRS Score)	1.96±0.59	3.16±0.45	2.18±0.75	3.29±0.61

The average immunoreactive score (IRS) of ALP expression in BMSCs group added with bovine DBM within normal culture was 3.20 ± 0.63 compared to 1.12 ± 0.36 in ASCs groups and had a statistically significant effect ($p = 0.000$). Thus in osteogenic culture, the IRS mean of the ALP expression in the BMSCs group compared to ASCs group added with bovine DBM also differ significantly ($p = 0.000$).




Figure 3. Number of osteoblast cell, immunoreactive score of ALP and osteocalcin for BMSCs and ASCs within normal and osteogenic culture, (*= $p < 0.05$ **= $p < 0.01$ ***= $p < 0.001$ #= $p > 0.05$)

The mean immunoreactive score of OC expression in the BMSCs group added with bovine DBM and cultured at normal condition was slightly lower than in the ASCs group (1.96 ± 0.59 vs 2.18 ± 0.75) but not statistically significant ($p = 0.498$). Whereas the average IRS of osteocalcin expression at osteogenic culture in BMSCs group was 3.16 ± 0.45 and 3.29 ± 0.61 in ASCs group but the results were not statistically significant ($p = 0.603$).




Figure 4. (A) HE staining on normal culture of BMSCs, (B) HE staining on normal culture of ASCs, (C) ALP expression in normal culture of BMSCs, (D) ALP expression in normal culture ASCs tissue, (E) OC expression in normal culture of BMSCs tissue, (F) OC expression result in normal culture ASCs tissue.




Figure 5. (A) HE staining on osteogenic culture of BMSCs, (B) HE Staining on osteogenic culture of ASCs (C) ALP expression in osteogenic culture of BMSCs, (D) ALP expression in osteogenic culture ASCs tissue, (E) OC expression in osteogenic culture of BMSCs, (F) OC expression result in osteogenic culture ASCs.

The use of stem cells in combination with scaffolds showed mixed results. The capability of mesenchymal stem cell differentiation (MSC) largely determines its use in managing various tissue damage. Stem cells derived from bone marrow are MSCs which was first studied and used widely. However, the morbidity induced when taking BMSCs causes other sources such as adipose and umbilical cord mesenchymal stem cells is further investigated [11]. Although most researchers conclude that BMSCs have better osteogenic proliferation capabilities than others [12-14], Kern et

al investigated the differentiation of bone marrow, adipose, and umbilical cord mesenchymal stem cells, became bone, and the results showed no significant difference between BMSCs and ASCs [15]. This led to the study using both types of stem cells growing at a rapid rate to handle orthopedic cases with bone defects.

Studies by Garmie et al and Janicki et al showed promising results from stem cell and scaffold applications of non-connective fractures [16,17]. However, this study supported by several other studies, shows that BMSCs remain superior to muscle, adipose and umbilical cord [18-20]. Other studies have suggested the use of embryonic stem cells with scaffolds with promising results, but limited cell resources and difficulty in processing require further study to obtain optimal results [21]. Liu et al used ASCs inserted into the scaffold of the shells to correct the cranial defects in experimental animals with satisfactory results [22].

Several experiments have been conducted to combine DBM with MSC. Schubert et al reported a significant difference between the use of ASCs implanted into DBM compared with the autopsy of the cancellous bones in spinal fusion using pig experiments [23]. Evaluations performed using CT scans and histologic examinations showed better performance in the group DBM added A-MSCs. Several other researchers have tried to combine DBM with BMSCs with satisfactory results [24,25].

The results of this study indicate a significant comparison in which the amount of osteoblasts and alkaline phosphatase expression produced by differentiation of BMSCs is higher than ASCs, but the resulting osteocalcin expression did not differ significantly. This is in line with the previous hypothesis that BMSCs have a higher osteogenic differentiation potential than ASCs. This study also proves that bovine DBM is able to function as a scaffold where stem cells differentiate into osteoblasts.

Studies using ASCs and the scaffold, both DBM and other scaffold forms, are still widely used to determine their efficacy in overcoming bone defects, as the use of BMSCs means increasing morbidity in patients in the form of pain in bone marrow collection sites. The authors argue that in vivo research is needed to directly compare the efficacy of DBM use in the treatment of bone defects in addition to ASCs or BMSCs.

4. Conclusion

Xenogenic bovine Demineralized Bone Matrix (DBM) is able to function as a scaffold where mesenchymal stem cells differentiate into osteoblast cells. Bone marrow mesenchymal stem cells have better osteogenic differentiation capabilities than adipose mesenchymal stem cells when implanted into Demineralized Bone Matrix.

References

- [1] Arun R. S., Michael C. M., Jeffrey O. H. Bone Tissue Engineering: State of the Union. Drug Discovery Today, 2014;19:781-786. <https://doi.org/10.1016/j.drudis.2014.04.010>
- [2] Natasha H., Matthew B. Focus on Management of Traumatic Bone Loss. Journal of Bone and Joint Surgery, 2011.
- [3] Greenwald AS, Boden SD, Goldberg VM, et al. 2002. Bone Graf Substitute: Facts, Fictions & Applications. Committee on Biological Implants. American Academy of Orthopaedic Surgeons. 69th Annual Meeting. Dallas, Texas.
- [4] Abdurrahman, 2002. Indonesian Tissue Banking: Progress on Tissue Production and Development of Quality Assurance, the 9th International Conference on Tissue Bank. Seoul.
- [5] Calori GM, Mazza E, Colombo M, Ripamonti C. The use of bone-graft substitutes in large bone defects: any specific needs? Injury, 2011 Sep;42 Suppl 2: S56-63.
- [6] Lementowski P.W. 2010. Acute and Chronic Complications of Intracortical Iliac Crest Bone Grafting Versus the Traditional Corticocancellous Technique for Spinal Fusion Surgery. Orthopedics, April 2010,33(4). <https://doi.org/10.3928/01477447-20100225-08>

-
- [7] Shibuya N1, Jupiter DC2Bone graft substitute: allograft and xenograft. Clin Podiatr Med Surg, 2015 Jan;32(1):21-34. <https://doi.org/10.1016/j.cpm.2014.09.011>
- [8] Sean M. Devitt et al. Successful Isolation of Viable Adipose-derived Stem Cells from Human Adipose Tissue Subject to Long-Term Cryopreservation: Positive Implications for Adult Stem Cell-Based Therapeutics in Patient of Advanced Age. Stem Cells International, 2015. <https://dx.doi.org/10.1155/2015/146421>
- [9] Sarvazyan, N. Cell and Tissue Engineering. Creation of Living Tissue: An Engineering Feat. Springer-Verlag Berlin Heidelberg. 2012, 1(8): 1-8.
- [10] Mahyudin F, Utomo DN, Suroto H, Wahyu T, Edward M, Gaol IL. Comparative Effectiveness of Bone Grafting using Xenograft Freeze-Dried Cortical Bovine, Allograft Freeze-Dried Cortical New Zealand White Rabbit, Xenograft Hydroxyapatite Bovine, and Xenograft Demineralized Bone Matrix Bovine in Bone Defect of Femoral. 2017;2017:1-11. <https://doi.org/10.1155/2017/7571523>
- [11] Arrington ED, Smith WJ, Chambers HG, et al. Complications of iliac crest bone graft harvesting. Clin Orthop Relat Res 1996;(329):300-9.
- [12] Bajada S, Mazakova I, Richardson JB, et al. Updates on stem cells and their applications in regenerative medicine. J Tissue Eng Regen Med 2008;2:169-83. <https://doi.org/10.1002/term.83>
- [13] Bianco P, Riminucci M, Gronthos S, et al. Bone marrow stromal stem cells: nature, biology, and potential applications. Stem Cells, 2001;19:180-92. <https://doi.org/10.1634/stemcells.19-3-180>
- [14] Niemeyer P, Kornacker M, Mehlhorn A, et al. Comparison of immunological properties of bone marrow stromal cells and adipose tissue-derived stem cells before and after osteogenic differentiation in vitro. Tissue Eng, 2007;13(1):111-21. <https://doi.org/10.1089/ten.2006.0114>
- [15] Kern, S et al. Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. Stem Cells, 2006;24(5):1294–1301. <https://doi.org/10.1634/stemcells.2005-0342>
- [16] Gamie, Z et al. Stem cells combined with bone graft substitutes in skeletal tissue engineering. Expert Opin Biol Ther, 2012;12(6):713-729. <https://doi.org/10.1517/14712598.2012.679652>
- [17] Janicki P, Schmidmaier G. What should be the characteristics of the ideal bone graft substitute? Combining scaffolds with growth factors and/or stem cells. Injury 2011;42:S77-81. <https://doi.org/10.1016/j.injury.2011.06.014>
- [18] Yoshimura H, Muneta T, Nimura A, et al. Comparison of rat mesenchymal stem cells derived from bone marrow, synovium, periosteum, adipose tissue, and muscle. Cell Tissue Res, 2007; 327: 449-62. <https://doi.org/10.1007/s00441-006-0308-z>
- [19] Schneider RK, Puellen A, Kramann R, et al. The osteogenic differentiation of adult bone marrow and perinatal umbilical mesenchymal stem cells and matrix remodeling in three-dimensional collagen scaffolds. Biomaterials, 2010;31:467-80. <https://doi.org/10.1016/j.biomaterials.2009.09.059>
- [20] Ohgushi H, Dohi Y, Tamai S, et al. Osteogenic differentiation of marrow stromal stem cells in porous hydroxyapatite ceramics. J Biomed Mater Res, 1993;27:1401-7.
- [21] Levenberg S, Huang NF, Lavik E, et al. Differentiation of human embryonic stem cells on three-dimensional polymer scaffolds. Proc Natl Acad Sci USA, 2003;100:12741-6. <https://doi.org/10.1073/pnas.1735463100>

- [22] Liu, G et al. 2013. Bone regeneration in a canine cranial model using allogeneic adipose derived stem cells and coral scaffold. *Biomaterials*, 2013;**34**(11):2655-2664. <https://doi.org/10.1016/j.biomaterials.2013.01.004>
- [23] Schubert, T et al. 2013. Critical size bone defect reconstruction by an autologous 3D osteogenic-like tissue derived from differentiated adipose MSCs. *Biomaterials*, 2013;**34**:4428-4438.
- [24] Mauney JR, Jaquiere C, Volloch V, et al. In vitro and in vivo evaluation of differentially demineralized cancellous bone scaffolds combined with human bone marrow stromal cells for tissue engineering. *Biomaterials*, 2005;**26**:3173-85.
<https://doi.org/10.1016/j.biomaterials.2004.08.020>
- [25] Liu G, Sun J, Li Y, et al. Evaluation of partially demineralized osteoporotic cancellous bone matrix combined with human bone marrow stromal cells for tissue engineering: an in vitro and in vivo study. *Calcif Tissue Int*, 2008;**83**:176-85. <https://doi.org/10.1007/s00223-008-9159-9>.

In Vitro Comparative Study of Osteogenic Differentiation Ability between Adipose and Bone Marrow Mesenchymal Stem Cell Applied to Bovine Demineralized Bone Matrix

ORIGINALITY REPORT

25%

SIMILARITY INDEX

15%

INTERNET SOURCES

23%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

www.iris.unina.it

Internet Source

1%

2

www.springerprotocols.com

Internet Source

1%

3

Ho, Angela W.H., and W.Y. Ip. "Biocompatibility and Efficacy of Five-Channel and Eight-Channel Crosslinked Urethane-Doped Polyester Elastomers (CUPEs) as Nerve Guidance Conduit for Reconstruction of Segmental Peripheral Nerve Defect Using Rat Model", Journal of Biomimetics Biomaterials and Biomedical Engineering, 2014.

Publication

1%

4

Ahmad Oryan, Amin Bigham-Sadegh, Somayeh Monazzah. "Characterization of turtle demineralized bone matrix (DBM) and turtle bone in rat radial bone defects; radiological, histopathological, biomechanical and scanning

1%

electron microscopic evaluation", Veterinarski arhiv, 2017

Publication

5	www.sistec.ac.in Internet Source	1 %
6	Somayeh Monazzah, Ahmad Oryan, Amin Bigham-Sadegh, Abdolhamid Meimandi-Parizi. "Application of bovine bone versus bovine DBM graft on bone healing of radial defect in rat", Comparative Clinical Pathology, 2017 Publication	1 %
7	www.stem-art.com Internet Source	1 %
8	dspace.ewha.ac.kr Internet Source	1 %
9	Gu, Yun Qing, Dong Shun Dai, Jie Gang Mou, Shui Hua Zheng, Deng Hao Wu, and Evan Wang. "Overview of the Technology of Bionic Surface Drag Reduction", Journal of Biomimetics Biomaterials and Biomedical Engineering, 2015. Publication	1 %
10	eprints.nottingham.ac.uk Internet Source	1 %
11	digituma.uma.pt Internet Source	1 %

12

www.ijbs.com

Internet Source

1 %

13

Ahmad Barghash, Nicole Golob-Schwarzl, Volkhard Helms, Johannes Haybaeck, Sonja M. Kessler. "Elevated expression of the *IGF2* mRNA binding protein 2 (IGF2BP2/IMP2) is linked to short survival and metastasis in esophageal adenocarcinoma", *Oncotarget*, 2016

Publication

1 %

14

Donald G. Phinney. "Isolation of Mesenchymal Stem Cells from Murine Bone Marrow by Immunodepletion", *Methods in Molecular Biology*TM, 2008

Publication

1 %

15

profdoc.um.ac.ir

Internet Source

1 %

16

"Bone and Cartilage Regeneration", Springer Nature, 2016

Publication

<1 %

17

www.thno.org

Internet Source

<1 %

18

Advanced Structured Materials, 2016.

Publication

<1 %

19

doaj.org

Internet Source

<1 %

20

Korkmaz-Icöz, Sevil, Samer Al Said, Tamás Radovits, Shiliang Li, Maik Brune, Péter Hegedűs, Ayhan Atmanli, Mihály Ruppert, Paige Brlecic, Lorenz Heyne Lehmann, Bernd Lahrmann, Niels Grabe, Yutaka Yoshikawa, Hiroyuki Yasui, Patrick Most, Matthias Karck, and Gábor Szabó. "Oral treatment with a zinc complex of acetylsalicylic acid prevents diabetic cardiomyopathy in a rat model of type-2 diabetes: activation of the Akt pathway", Cardiovascular Diabetology, 2016.

Publication

<1 %

21

yournewswire.com

Internet Source

<1 %

22

d-nb.info

Internet Source

<1 %

23

Peng, Xuewu Song, Tongxing Hu, Xiaoming . "Phenotypic and functional properties of porcine dedifferentiated fat cells during the long-term cult", BioMed Research International, Annual 2015 Issue

Publication

<1 %

24

Yaling Shi. "Adipose Derived Stem Cells Combined with a Demineralized Cancellous Bone Substrate for Bone Regeneration", Tissue Engineering Part A, 04/13/2012

Publication

<1 %

25

opus.bibliothek.uni-wuerzburg.de

Internet Source

<1 %

26

benthamopen.com

Internet Source

<1 %

27

Yu-Jen Chang, Daniel Tzu-bi Shih, Ching-Ping Tseng, Tzu-Bou Hsieh, Don-Ching Lee, Shiaw-Min Hwang. "Disparate Mesenchyme-Lineage Tendencies in Mesenchymal Stem Cells from Human Bone Marrow and Umbilical Cord Blood", Stem Cells, 2006

Publication

<1 %

28

A. A. Ismail. "Effects of High-Mobility Group A Protein Application on Canine Adipose-Derived Mesenchymal Stem Cells In Vitro", Veterinary Medicine International, 2012

Publication

<1 %

29

www.dovepress.com

Internet Source

<1 %

30

oro.open.ac.uk

Internet Source

<1 %

31

stemcellres.com

Internet Source

<1 %

32

mdpi.com

Internet Source

<1 %

33

Paul S. Pipitone, Saqib Rehman. "Management

of Traumatic Bone Loss in the Lower Extremity",
Orthopedic Clinics of North America, 2014

Publication

<1 %

34

ecmjournal.org

Internet Source

<1 %

35

Dequan Kong, Guoqiang Su, Lang Zha, Hongyu Zhang, Jifeng Xiang, Wei Xu, Yucheng Tang, Ziwei Wang. "Coexpression of HMGA2 and Oct4 predicts an unfavorable prognosis in human gastric cancer", Medical Oncology, 2014

Publication

<1 %

36

H Aurich. "Hepatocyte differentiation of mesenchymal stem cells from human adipose tissue in vitro promotes hepatic integration in vivo", Gut, 04/01/2009

Publication

<1 %

37

jacobspublishers.com

Internet Source

<1 %

38

www.tdx.cat

Internet Source

<1 %

39

eprints.unife.it

Internet Source

<1 %

40

Tommasi, Giacomo, Stefano Perni, and Polina Prokopovich. "An Injectable Hydrogel as Bone Graft Material with Added Antimicrobial Properties", Tissue Engineering Part A, 2016.

<1 %

41	P Niemeyer. "Transplantation of mesenchymal stromal cells on mineralized collagen leads to ectopic matrix synthesis in vivo independently from prior in vitro differentiation", Cytotherapy, 8/1/2006	<1 %
----	---	------

Publication

42	edoc.unibas.ch	<1 %
----	--	------

Internet Source

43	Perinatal Stem Cells, 2014.	<1 %
----	-----------------------------	------

Publication

44	stemcells.alphamedpress.org	<1 %
----	--	------

Internet Source

45	www.faqs.org	<1 %
----	--	------

Internet Source

46	imo2.thejakartapost.com	<1 %
----	--	------

Internet Source

47	Kelly A. Lenton. "Re: Porous Polymethylmethacrylate as Bone Substitute in the Craniofacial Area.", Journal of Craniofacial Surgery, 07/2003	<1 %
----	---	------

Publication

48	www.squeezedfatgrafting.com	<1 %
----	--	------

Internet Source

49	Hu, Xiaoqing, Xin Zhang, Linghui Dai, Jingxian	
----	--	--

Zhu, Zhuqing Jia, Weiping Wang, Chunyan Zhou, and Yingfang Ao. "Histone Deacetylase Inhibitor Trichostatin A Promotes the Osteogenic Differentiation of Rat Adipose-Derived Stem Cells by Altering the Epigenetic Modifications on *Runx2* Promoter in a BMP Signaling-Dependent Manner", *Stem Cells and Development*, 2013.

Publication

<1 %

50

I. Sekiya. "Synovial fluid-derived mesenchymal stem cells increase after intra-articular ligament injury in humans", *Rheumatology*, 04/29/2008

Publication

<1 %

51

Mandi J. Lopez. "Acceleration of spinal fusion using syngeneic and allogeneic adult adipose derived stem cells in a rat model", *Journal of Orthopaedic Research*, 03/2009

Publication

<1 %

52

Westhauser, F., C. Weis, M. Hoellig, T. Swing, G. Schmidmaier, M.-A. Weber, W. Stiller, H.-U. Kauczor, and A. Moghaddam. "Heidelberg-mCT-Analyzer: a novel method for standardized microcomputed-tomography-guided evaluation of scaffold properties in bone and tissue research", *Royal Society Open Science*, 2015.

Publication

<1 %

53

Tropel, P.. "Isolation and characterisation of

<1 %

mesenchymal stem cells from adult mouse bone marrow", Experimental Cell Research, 20040501

Publication

54

Yongsoo Choi. "A Bridging Demineralized Bone Implant Facilitates Posterolateral Lumbar Fusion in New Zealand White Rabbits", Spine, 01/2007

Publication

55

Lementowski, Peter W., Paul Lucas, and Rudolph F. Taddonio. "Acute and Chronic Complications of Intracortical Iliac Crest Bone Grafting Versus the Traditional Corticocancellous Technique for Spinal Fusion Surgery", Orthopedics, 2010.

Publication

56

Patricia A. Zuk. "Multilineage Cells from Human Adipose Tissue: Implications for Cell-Based Therapies", Tissue Engineering, 04/2001

Publication

57

Dwikora N. Utomo, Ferdiansyah Mahyudin, Kukuh D. Hernugrahanto, Heri Suroto, Muhammad Zaim Chilmi, Fedik Abdul Rantam. "Implantation of platelet rich fibrin and allogenic mesenchymal stem cells facilitate the healing of muscle injury: An experimental study on animal", International Journal of Surgery Open, 2018

Publication

<1 %

<1 %

<1 %

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

In Vitro Comparative Study of Osteogenic Differentiation Ability between Adipose and Bone Marrow Mesenchymal Stem Cell Applied to Bovine Demineralized Bone Matrix

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8